PHYSICIAN'S REPORT—CHILD CARE CENTERS

(CHILD'S PRE-ADMISSION HEALTH EVALUATION)

PART A – PARENT'S CONSENT (TO BE COMPLETED BY PARENT)

(NAME OF CHILD)

__, born __

(BIRTH DATE)

_ is being studied for readiness to enter

_. This Child Care Center/School provides a program which extends from _____: ____

(NAME OF CHILD CARE CENTER/SCHOOL)

a.m./p.m. to ______ a.m./p.m. , _____ days a week.

Please provide a report on above-named child using the form below. I hereby authorize release of medical information contained in this report to the above-named Child Care Center.

(SIGNATURE OF PARENT, GUARDIAN, OR CHILD'S AUTHORIZED REPRESENTATIVE)

(TODAY'S DATE)

PART B - PHYSICIAN'S REPORT (TO BE COMPLETED BY PHYSICIAN)

Problems of which you should be aware:	
Hearing:	Allergies: medicine:
с. С	0
Vision:	Insect stings:
	nood dange.
Developmental:	Food:
Developmental.	1000.
	A 11
Language/Speech:	Asthma:
Dental:	
Other (Include behavioral concerns):	
Comments/Explanations:	

MEDICATION PRESCRIBED/SPECIAL ROUTINES/RESTRICTIONS FOR THIS CHILD:

IMMUNIZATION HISTORY: (Fill out or enclose California Immunization Record, PM-298.)

VACCINE	DATE EACH DOSE WAS GIVEN						
	1st	2nd	3rd	4th	5th		
POLIO (OPV OR IPV)	/ /	/ /	/ /	/ /	/ /		
DTP/DTaP/ (DIPHTHERIA, TETANUS AND [ACELLULAR] PERTUSSIS OR TETANUS DT/Td AND DIPHTHERIA ONLY)	/ /	/ /	/ /	/ /	/ /		
MMR (MEASLES, MUMPS, AND RUBELLA)	/ /	/ /		· · · ·			
(REQUIRED FOR CHILD CARE ONLY) HIB MENINGITIS (HAEMOPHILUS B)		/ /	/ /	/ /			
HEPATITIS B	/ /	/ /	/ /				
VARICELLA (CHICKENPOX)	/ /	/ /					
SCREENING OF TB RISK FACTOR		· · ·					
Risk factors present; Manto previous positive skin test d Communicable TB dise	ocumented).	ormed (unless					
I have have not	reviewed the	above information w	vith the parent/guar	dian.			
Address: Date		Date	This Form Complete	ed:			
		F	hysician 🗹 Pl	nysician's Assistant	✓ Nurse Practitioner		

RISK FACTORS FOR TB IN CHILDREN:

- * Have a family member or contacts with a history of confirmed or suspected TB.
- * Are in foreign-born families and from high-prevalence countries (Asia, Africa, Central and South America).
- * Live in out-of-home placements.
- * Have, or are suspected to have, HIV infection.
- * Live with an adult with HIV seropositivity.
- * Live with an adult who has been incarcerated in the last five years.
- * Live among, or are frequently exposed to, individuals who are homeless, migrant farm workers, users of street drugs, or residents in nursing homes.
- * Have abnormalities on chest X-ray suggestive of TB.
- * Have clinical evidence of TB.

Consult with your local health department's TB control program on any aspects of TB prevention and treatment.